

1/(N - 1) expansion approach to full counting statistics for the SU(N) Anderson model

Akira Oguri

Department of Physics, Osaka City University, Sumiyoshi-ku, Osaka, Japan

Rui Sakano

Institute for Solid State Physics, University of Tokyo, Kashiwa, Chiba, Japan

(Dated: October 12, 2012)

We apply a recently developed $1/(N - 1)$ expansion to the full counting statistics for the N -fold degenerate Anderson impurity model in the Kondo regime. This approach is based on the perturbation theory in the Coulomb interaction U , and is different from the conventional large N theories such as the usual $1/N$ expansion and non-crossing approximation. We have confirmed that the calculations carried out up to order $1/(N - 1)^2$ agree closely with those of the numerical renormalization group at $N = 4$, where the degeneracy is still not so large. This ensures the applicability of our approach for $N \geq 4$. We present the results of the cumulants of the probability distribution function for nonequilibrium current through a quantum dot in the particle-hole symmetric case.

PACS numbers: 72.15.Qm, 73.63.Kv, 75.20.Hr

Keywords: Kondo effect, quantum dot, orbital degeneracy, nonequilibrium current

I. INTRODUCTION

The universal Kondo behavior in the quantum dots has been studied intensively for nonequilibrium current [1–6] and shot noise [7–13]. Effects of orbital degeneracy on the universality have also been a subject of current interest [14–18].

Essential features of the low-energy properties of the orbital systems may be deduced from the SU(N) Anderson model, which for $N = 2$ corresponds to the spin degeneracy. The exact numerical renormalization group (NRG) approach [19] has been applied to this model for small degeneracies $N \leq 4$ [20]. For $N > 4$, the conventional large N theories, such as the $1/N$ expansion and the non-crossing approximation [21, 22] are applicable to a wide energy scale except for the low-energy Fermi-liquid regime. Therefore, at low energies alternative approaches are needed to explore the universal behavior for $N > 4$.

Recently, we proposed a different type of large N theory, using a scaling that takes $u = (N - 1)U$ as an independent variable instead of the bare Coulomb interaction U [23, 24]. The factor $N - 1$ represents the number of interacting orbitals, excluding the one prohibited by the Pauli principle. With this scaling the perturbation series in U can be reorganized as an expansion in powers of $1/(N - 1)$. To leading order in $1/(N - 1)$ it describes the Hartree-Fock random phase approximation (HF-RPA). The higher-order corrections systematically describe the fluctuations beyond the HF-RPA. As the unperturbed Hamiltonian includes the tunneling matrix element between the impurity and conduction bands, this approach naturally describes the Fermi-liquid state.

In this report, we discuss dependence of the renormalized parameters on the degeneracy N , carrying out the calculations up to order $1/(N - 1)^2$. Furthermore, we apply this approach to the full counting statistics [25–27] for the nonequilibrium current distribution through quantum dots in the Kondo regime.

II. MODEL AND FORMULATION

We consider the SU(N) impurity Anderson model with a finite interaction U , connected to two leads ($\nu = L, R$): $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_U$,

$$\mathcal{H}_0 = \sum_{m=1}^N \xi_d d_m^\dagger d_m + \sum_{\nu=L,R} \sum_{m=1}^N \int_{-D}^D d\epsilon \epsilon c_{\epsilon\nu m}^\dagger c_{\epsilon\nu m} + \sum_{\nu=L,R} \sum_{m=1}^N v_\nu \left(d_m^\dagger \psi_{\nu m} + \psi_{\nu m}^\dagger d_m \right), \quad (1)$$

$$\mathcal{H}_U = \sum_{m \neq m'} \frac{U}{2} \left(n_{dm} - \frac{1}{2} \right) \left(n_{dm'} - \frac{1}{2} \right). \quad (2)$$

Here, $\xi_d = \epsilon_d + (N - 1)U/2$, $d_m^\dagger$ is the creation operator for an electron with energy ϵ_d and orbital m ($= 1, 2, \dots, N$) in the impurity site, and $n_{dm} = d_m^\dagger d_m$. The operator $c_{\epsilon\nu m}^\dagger$ for a conduction electron in the lead ν is normalized as $\{c_{\epsilon\nu m}, c_{\epsilon'\nu' m'}^\dagger\} = \delta_{\nu\nu'} \delta_{mm'} \delta(\epsilon - \epsilon')$, and $\psi_{\nu m} = \int_{-D}^D d\epsilon \sqrt{\rho} c_{\epsilon\nu m}$. The hybridization energy scale is given by $\Delta \equiv \Gamma_L + \Gamma_R$, with $\Gamma_\nu = \pi \rho v_\nu^2$ and $\rho = 1/(2D)$.

We use the imaginary-frequency Green's function that is given, for $|\omega| \ll D$, by

$$G(i\omega) = \frac{1}{i\omega - \xi_d + i\Delta \operatorname{sgn} \omega - \Sigma(i\omega)}. \quad (3)$$

Here, $\Sigma(i\omega)$ is the self-energy due to $\mathcal{H}_U$. The ground-state average of the local charge, $\langle n_{dm} \rangle = \delta/\pi$, can be deduced from the phase shift $\delta \equiv \cot^{-1}(E_d^*/\Delta)$ with $E_d^* \equiv \xi_d + \Sigma(0)$. The renormalized parameters are defined by $1/z \equiv 1 - \partial \Sigma(i\omega)/\partial(i\omega)|_{\omega=0}$, $\tilde{\epsilon}_d \equiv z E_d^*$, and $\tilde{\Delta} \equiv z \Delta$. Furthermore, the enhancement factor for the spin and charge susceptibilities, $\tilde{\chi}_s \equiv \tilde{\chi}_{mm} - \tilde{\chi}_{mm'}$ and $\tilde{\chi}_c \equiv \tilde{\chi}_{mm} + (N - 1)\tilde{\chi}_{mm'}$, can be expressed in terms of z and the vertex function $\Gamma_{mm';m'm}(i\omega_1, i\omega_2; i\omega_3, i\omega_4)$

$$\text{wavy line} = \text{wavy line} + \text{bubble} + \text{bubble} + \dots$$

FIG. 1: The leading order diagrams in the $1/(N-1)$ expansion. The wavy and solid lines indicate U and G_0 . The double wavy line represents $\mathcal{U}_{\text{bub}}(i\omega)$ given in Eq. (7).



FIG. 2: The order $1/(N-1)$ vertex and self-energy.

for $m \neq m'$ [28, 29]:

$$\tilde{\chi}_{mm} = \frac{1}{z}, \quad \tilde{\chi}_{mm'} = -\frac{\sin^2 \delta}{\pi \Delta} \Gamma_{mm';m'm}(0,0;0,0), \quad (4)$$

and $\tilde{U} \equiv z^2 \Gamma_{mm';m'm}(0,0;0,0)$ for $m \neq m'$ represents the residual interaction between the quasi-particles.

We introduce a scaling for the bare and renormalized interactions, by multiplying a factor $N-1$ [23, 24],

$$g \equiv \frac{(N-1)U}{\pi \Delta}, \quad \tilde{g} \equiv \frac{(N-1)\tilde{U}}{\pi \tilde{\Delta}}. \quad (5)$$

With these parameters, the Wilson ratio $R \equiv z \tilde{\chi}_s$ and that for the charge sector can be expressed in the form,

$$R = 1 + \frac{\tilde{g}}{N-1} \sin^2 \delta, \quad z \tilde{\chi}_c = 1 - \tilde{g} \sin^2 \delta. \quad (6)$$

One of the merit of this scaling is that the perturbation expansion with respect to $\mathcal{H}_U$ can be classified according to the power of $1/(N-1)$, taking g as an independent variable. In the present report we consider the particle-hole symmetric case: $\epsilon_d = -(N-1)U/2$ and $\delta = \pi/2$.

III. EXPANSION UP TO ORDER $1/(N-1)^2$

At the zero order with respect to $1/(N-1)$, the limit of $N \rightarrow \infty$ is taken at fixed g , and it gives the Hartree-Fock results. Specifically, at half-filling the unperturbed Green's function is given by $G_0(i\omega) = [i\omega + i\Delta \text{sgn} \omega]^{-1}$ as $\xi_d = 0$.

The leading order corrections in the $1/(N-1)$ expansion arise from a series of the bubble diagrams, $\mathcal{U}_{\text{bub}}(i\omega)$, of the RPA type shown in Fig. 1,

$$\mathcal{U}_{\text{bub}}(i\omega) = U + U \sum_{k=1}^{\infty} \mathcal{A}_k \left[-U \chi_0(i\omega) \right]^k. \quad (7)$$

Here, $\chi_0(i\omega) = -\int \frac{d\omega'}{2\pi} G_0(i\omega + i\omega') G_0(i\omega')$, and the coefficient $\mathcal{A}_k = (N-1)^k \sum_{p=0}^k [-1/(N-1)]^p$ arises from the summations over the orbital indices for a series of k fermion loops. Correspondingly, the order $1/(N-1)$ contributions of $\Gamma_{mm';m'm}(0,0;0,0)$ arise from the first




FIG. 3: The order $1/(N-1)^2$ diagrams for the vertex function $\Gamma_{mm';m'm}(0,0;0,0)$ for $m \neq m'$ at half-filling $\xi_d = 0$.




FIG. 4: The order $1/(N-1)^2$ self-energy at half-filling.

diagram in Fig. 2, which give the leading order correction to the renormalized coupling,

$$\tilde{g} = \frac{g}{1+g} + O\left(\frac{1}{N-1}\right). \quad (8)$$

This correction determines the Wilson ratio R to order $1/(N-1)$ through Eq. (6). Similarly, the order $1/(N-1)$ self-energy arises from the second diagram in Fig. 2.

Higher order fluctuations beyond the RPA appear first through the next-leading-order contributions. Figures 3 and 4 show the order $1/(N-1)^2$ diagrams for the vertex function and self-energy in the particle-hole symmetric case, respectively. These contributions and the higher order components from Eq. (7) determine the renormalization factor z and the Wilson ratio R to order $1/(N-1)^2$. In the present work, we have calculated all these next-leading-order contributions. The outline of the calculations were described in a previous paper [24]. We have also carried out the NRG calculations for $N = 4$ to demonstrate the reliability of the $1/(N-1)$ expansion.

Figure 5 shows the next-leading-order results for $\tilde{g}$ and z , plotted as a function of g . We see the very close agreement between the NRG and the next-leading-order results in the $1/(N-1)$ expansion for $N = 4$, especially for the renormalized coupling $\tilde{g}$. The two curves for $\tilde{g}$, for $N = 4$, almost overlap each other over the whole range of g although the next-leading-order results (green dashed line) are slightly smaller than those of the NRG (solid circles). As N increases, $\tilde{g}$ converges rapidly to the RPA value $\tilde{g} \rightarrow g/(1+g)$, which is asymptotically exact in the limit of $N \rightarrow \infty$. We also see that the value that $\tilde{g}$ can take is bounded in a very narrow region between the curve for $N = 4$ and that for the $N \rightarrow \infty$ limit.

The order $1/(N-1)^2$ results for the renormalization factor z , shown in Fig. 5, also agree well with the NRG




FIG. 5: (Color online) Renormalized parameters $\tilde{g}$ and z in the particle-hole symmetric case $\epsilon_d = -(N-1)U/2$ are plotted as a function of the scaled Coulomb interaction g for $N = 2$ (Bethe ansatz [30]), and for $N = 4, 6, 8$ (next-leading order in the $1/(N-1)$ expansion). For $N = 4$, the NRG results [16, 31] are also shown with the solid circle ($\bullet$). In the $N \rightarrow \infty$ limit, $\tilde{g}$ and z approach $\tilde{g} \rightarrow g/(1+g)$ and $z \rightarrow 1.0$.

results for $N = 4$ at $g \lesssim 3.0$, or equivalently in the region where $\tilde{g} \lesssim 0.8$. This indicates that the next-leading-order results also reproduce the Kondo energy scale, $\tilde{\Delta} = z\Delta$, properly from the weak-coupling to intermediate-coupling regions where $\tilde{g}$ is still not very close to 1.0, the value in the strong coupling limit.

IV. FULL COUNTING STATISTICS

The $1/(N-1)$ expansion can be applied to nonequilibrium transport at finite bias voltages V . To be specific, we choose the lead-dot couplings and chemical potentials in the leads to be symmetric $\Gamma_L = \Gamma_R$ and $\mu_L = -\mu_R (= eV/2)$, and consider a steady state in the particle-hole symmetric case. In this case, an asymptotically exact result was obtained for the retarded Green's function at low energies up to terms of order ω^2 , T^2 , and $(eV)^2$ [4, 16],

$$G^r(\omega) \simeq \frac{z}{\omega + i\tilde{\Delta} + i\frac{\tilde{g}^2}{2(N-1)\Delta} [\omega^2 + \frac{3}{4}(eV)^2 + (\pi T)^2]}. \quad (9)$$

The average and fluctuations of the steady current at low energies can be described by the local Fermi-liquid theory, which follows from this form of the Green's function. The Fermi-liquid theory, or the related renormalized perturbation theory (RPT) [32], can also be applied to the full counting statistics for nonequilibrium transport.

Specifically, we consider the probability distribution $P(\mathbf{q})$ of the transferred charge $\mathbf{q} = (q_1, q_2, \dots, q_N)$ from the left lead to the orbital m of the Anderson impurity during a time interval $\mathcal{T}$. The generating function for this probability distribution is defined by $\chi(\boldsymbol{\lambda}) = \sum_{\mathbf{q}} e^{i\boldsymbol{\lambda} \cdot \mathbf{q}} P(\mathbf{q})$ with the counting fields $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_N)$, and can be expressed in the form [25],

$$\chi(\boldsymbol{\lambda}) = \left\langle T_C \exp \left\{ -i \int_C dt [\mathcal{H}_T^\lambda(t) + \mathcal{H}_U(t)] \right\} \right\rangle. \quad (10)$$

Here, T_C is the time ordering operator along the Keldysh contour C , and the time evolution is defined with respect to an extended Hamiltonian $\mathcal{H}^\lambda(t) = \mathcal{H}_0 + \mathcal{H}_T^\lambda(t) + \mathcal{H}_U$,

$$\mathcal{H}_T^\lambda(t) = \sum_{km} \left[v_L e^{i\lambda_m(t)/2} d_m^\dagger c_{kLm} + v_R d_m^\dagger c_{kRm} \right] + \text{H.c.} \quad (11)$$

Here, the counting field depends on the path such that $\lambda_m(t_\mp) = \pm \lambda_m$ for the forward (t_-) and backward (t_+) paths, respectively, and is switched on during a time interval $[t: 0 \rightarrow \mathcal{T} \rightarrow 0]$.

We obtained an explicit expression of $\chi(\boldsymbol{\lambda})$, which is asymptotically exact at low energies up to order $(eV)^3$ for general N and U , using the RPT [17, 18]. The cumulant for the full current can be derived from $\chi(\boldsymbol{\lambda})$, choosing the counting fields to be $\lambda_m = \lambda$ for all m , and then taking a derivative $\mathcal{C}_n = (-i)^n \frac{d^n}{d\lambda^n} \ln \chi(\lambda)$:

$$\frac{\mathcal{C}_n}{\mathcal{T}} = \frac{J_u}{e} \left[\delta_{1n} + \frac{(-1)^n}{12} \left\{ 1 + \frac{(1+2^{n+1})\tilde{g}^2}{(N-1)} \right\} \left(\frac{eV}{\tilde{\Delta}} \right)^2 \right]. \quad (12)$$

Here, $J_u = Ne^2V/(2\pi\hbar)$ is the linear-response current of the unitary limit, and $\delta_{nn'}$ is the Kronecker's delta. Equation (12) shows that the nonequilibrium properties are also characterized by the two renormalized parameters, $\tilde{g}$ and the Kondo energy scale $\tilde{\Delta}$, at low energies in the particle-hole symmetric case. For $n = 1$, the cumulant $e\mathcal{C}_1/\mathcal{T}$ expresses the steady current. The universal properties of the cumulants for $n \geq 2$ can be extracted from the Fano-factor *inspired* ratio (FFIR), which is normalized with respect to the backscattering current $J_b \equiv J_u - e\mathcal{C}_1/\mathcal{T}$ [10]:

$$\frac{\mathcal{C}_n}{\mathcal{C}_n^P} = \frac{1 + \frac{(1+2^{n+1})\tilde{g}^2}{N-1}}{1 + \frac{5\tilde{g}^2}{N-1}}, \quad \frac{\mathcal{C}_n^P}{\mathcal{T}} \equiv (-1)^n \frac{J_b}{e}. \quad (13)$$

Here, $\mathcal{C}_n^P$ can also be regarded as the Poisson value of the cumulant, and the FFIR for $n = 2$ corresponds to the Fano factor [16].

Figure 6 shows the ratio $\mathcal{C}_n/\mathcal{C}_n^P$ for the noise $n = 2$ (top), the skewness $n = 3$ (middle), and sharpness $n = 4$ (bottom) as a function of the scaled Coulomb interaction g for several value of the orbital degeneracy N .




FIG. 6: (Color online) The FFIR C_n/C_n^P for $n = 2$ (top), $n = 3$ (middle), and $n=4$ (bottom) are plotted as a function of g for $N = 2$ (Bethe ansatz), and $N = 4, 6, 8, 10, 20$ (next-leading-order results in the $1/(N-1)$ expansion). The NRG results (solid circle) are also shown for $N = 4$.

We see the close agreement between the NRG and the next-leading-order results for $N = 4$ again. As the next-leading-order $1/(N-1)$ results for $\tilde{g}$ are numerically almost exact for $N \geq 4$, the results shown in Fig. 6 capture orbital effects correctly. In the strong coupling limit $g \rightarrow \infty$, the FFIR approach $C_n/C_n^P \rightarrow (N+2^{n+1})/(N+4)$ as the renormalized coupling converges to $\tilde{g} \rightarrow 1$. As N increases, $\tilde{g}$ converges rapidly to the value, $\tilde{g} \simeq g/(1+g)$, for the large N limit, as mentioned in the above. Thus, for $N \gtrsim 8$, the N dependence is determined essentially by the factor $1/(N-1)$, appearing explicitly in Eq. (13). In the limit of $N \rightarrow \infty$, the cumulant approaches the Poisson value, $C_n/C_n^P \rightarrow 1$, as the fluctuations due to electron correlations are suppressed and the mean-field theory becomes asymptotically exact.

In summary, carrying out the $1/(N-1)$ expansion up to order $1/(N-1)^2$, we have obtained the cumulants for the probability distribution of the current through a quantum dot with the orbital degeneracy $N > 4$ at low energies. The results of the renormalized coupling $\tilde{g}$ show the excellent agreement with the exact NRG results already at $N = 4$ in the particle-hole symmetric case. This enable us to obtain *almost* exact numerical results for the Fano-factor inspired ratio C_n/C_n^P for $N > 4$.

Acknowledgments

The authors thank T. Fujii, T. Kato, and A. C. Hewson for discussions. This work is supported by the JSPS Grant-in-Aid for Scientific Research C (No. 23540375, and No. 24540316). Numerical computation was partly carried out at the Yukawa Institute Computer Facility.

-
- [1] M. Grobis, I. G. Rau, R. M. Potok, H. Shtrikman, and D. Goldhaber-Gordon, Phys. Rev. Lett. **100**, 246601 (2008).
 - [2] G. D. Scott, Z. K. Keane, J. W. Ciszek, J. M. Tour, and D. Natelson, Phys. Rev. B **79**, 165413 (2009).
 - [3] A. Kaminski, Yu. V. Nazarov, and L. I. Glazman, Phys. Rev. B **62**, 8154 (2000).
 - [4] A. Oguri, Phys. Rev. B **64**, 153305 (2001).
 - [5] T. Fujii and K. Ueda, Phys. Rev. B **68**, 155310 (2003).
 - [6] A. C. Hewson, J. Bauer, and A. Oguri, J. Phys.: Condes. Matter. **17**, 5413 (2005).
 - [7] O. Zarchin, M. Zaffalon, M. Heiblum, D. Mahalu, and V. Umansky, Phys. Rev. B **77**, 241303 (2008).
 - [8] Y. Yamauchi, *et al.*, Phys. Rev. Lett. **106**, 17660 (2011).
 - [9] A. O. Gogolin and A. Komnik, Phys. Rev. B **73**, 195301 (2006).
 - [10] A. O. Gogolin, and A. Komnik, Phys. Rev. Lett. **97**, 016602 (2006).
 - [11] A. Golub, Phys. Rev. B **73**, 233310 (2006).
 - [12] E. Sela and J. Malecki, Phys. Rev. B **80**, 233103 (2009).
 - [13] T. Fujii, J. Phys. Soc. Jpn. **79**, 044714 (2010).
 - [14] T. Delattre, *et al.*, Nature Phys. **5**, 208 (2009).
 - [15] C. Mora, P. Vitushinsky, X. Leyronas, A. A. Clerk, and K. Le Hur, Phys. Rev. B **80**, 155322 (2009).
 - [16] R. Sakano, T. Fujii, and A. Oguri, Phys. Rev. B **83**, 075440 (2011).
 - [17] R. Sakano, A. Oguri, T. Kato and S. Tarucha, Phys. Rev. B **83**, 241301 (2011).
 - [18] R. Sakano, Y. Nishikawa, A. Oguri, A. C. Hewson, and S. Tarucha, Phys. Rev. Lett. **108**, 266401 (2012).
 - [19] H. R. Krishna-murthy, J. W. Wilkins, and K. G. Wilson, Phys. Rev. B **21**, 1003 (1980).
 - [20] W. Izumida, O. Sakai and Y. Shimizu, J. Phys. Soc. Jpn. **67** 2444 (1998).
 - [21] P. Coleman, Phys. Rev. B **28**, 5255 (1983).
 - [22] N. Bickers, Rev. Mod. Phys. **59**, 845 (1987).
 - [23] A. Oguri, R. Sakano, and T. Fujii, Phys. Rev. B **84**, 113301 (2011).
 - [24] A. Oguri, Phys. Rev. B **85**, 155404 (2012).
 - [25] L. S. Levitov, and M. Reznikov, Phys. Rev. B **70**, 115305 (2004).
 - [26] D. Bagrets, Y. Utsumi, D. Golubev, and G. Schön, Fortschr. Phys. **54**, 917 (2006).
 - [27] M. Esposito, U. Harbola, and S. Mukamel, Rev. Mod. Phys. **81**, 1665 (2009).
 - [28] K. Yamada, Prog. Theor. Phys. **53**, 970 (1975).
 - [29] A. Yoshimori, Prog. Theor. Phys. **55**, 67 (1976).
 - [30] V. Zlatić and B. Horvatić, Phys. Rev. B **28**, 6904 (1983).
 - [31] Y. Nishikawa, D. J. G. Crow, and A. C. Hewson, Phys. Rev. B **82**, 115123 (2010).
 - [32] A. C. Hewson, J. Phys. Condes. Matter **13**, 10011 (2001).